

Herpetological Review

Volume 47, Number 1 – March 2016



mabouia colonies on Turks and Caicos Bank islands (Reynolds and Niemiller 2009. *Herpetol. Rev.* 40:452; Powell and Henderson 2012, *op. cit.*). With its occupation of Great Inagua, this exotic species has established itself within a minimum of 16 Bahamian islands on five of the 15 island banks (Powell and Henderson 2012, *op. cit.*).

AARON H. GRIFFING (e-mail: agriffing66@gmail.com) and **AARON M. BAUER**, Department of Biology, Villanova University, 800 Lancaster Avenue, Villanova, Pennsylvania 19085, USA (e-mail: aaron.bauer@villanova.edu).

HEMIDACTYLUS PARVIMACULATUS (Sri Lankan Spotted House Gecko). USA: LOUISIANA: St. TAMMANY PARISH: private property ca. 4 km S of Abita Springs, E of State Hwy 59, and N of Interstate 12 (30.44000°N, 90.02000°W; WGS 84). 18 August 2013. Brad M. Glorioso. Verified by David Heckard. Florida Museum of Natural History (UF 176422, photo voucher). New parish record. This species was first reported in the Americas in the vicinity of Audubon Zoo in New Orleans, Louisiana (Heckard et al. 2013. *IRCF Reptiles & Amphibians* 20:192–196). This is the third report of this species in Louisiana (Heckard et al. 2013, *op. cit.*; Borgardt 2015. *Herpetol. Rev.* 46:217), and is now documented from Orleans, Jefferson, and St. Tammany parishes in southeast Louisiana. The individual was located at night, ca. 10 m from a dwelling on the forest floor amid a downed picket fence, which was resting atop a thick layer of pine needles. There are *H. turcicus* at the property, but this individual was recognized as unusual, and many photos were taken before releasing the animal. It was not until much later that it was determined to be *H. parvimaculatus*. The origin of this individual is unknown, as the owners of this 1.62-ha property are elderly and do not keep any pets. Subsequent casual searches have not turned up any new individuals. I thank David Heckard for his help with identification and discussion of this species in Louisiana.

BRAD M. GLORIOSO, U.S. Geological Survey, Wetland and Aquatic Research Center, Lafayette, Louisiana 70506, USA, e-mail: gloriosob@usgs.gov.

HEMIDACTYLUS TURCICUS (Mediterranean Gecko). USA: FLORIDA: SANTA ROSA Co.: Gulf Breeze, Reserve Blvd. at intersection with Heronwalk Dr. (30.40780°N, 86.95524°W; WGS 84). 7 October 2015. Ashton den Dekker. Auburn University Natural History Museum (AUM AHAP-D 1133, photo voucher). New county record (Krysko et al. 2011. *Atlas of Amphibians and Reptiles in Florida*. Final report, Florida Fish and Wildlife Conservation Commission, Tallahassee, Florida. 524 pp.). One of several individuals frequently observed at windows and exterior lights of private residence. The nearest previously documented location in the state occurs 20 km to the west at Pensacola Beach, Escambia County. This record fills a gap in the western Florida Panhandle. Thanks to David Laurencio for verifying the identification and Kevin Enge for confirming the county record. Additional records were examined for county occurrences via VertNet and Zoological Record.

BRIAN D. HOLT, Alabama Department of Conservation and Natural Resources, State Lands Division, Natural Heritage Section, 64 N Union Street, Suite 464, Montgomery, Alabama 36130, USA (e-mail: brian.holt@dcnr.alabama.gov); **ASHTON DEN DEKKER**, 2008 Reserve Blvd., Gulf Breeze, Florida 32563, USA.

HEMIDACTYLUS TURCICUS (Mediterranean Gecko). USA: TEXAS: BASTROP Co.: Redfield Farm (2158 FM 812), TX FM 812 ca.

1.3 rd km SE jct TX Hwy 21 (30.06377°N, 97.56084°W; WGS 84). 6 August 2014. Drew R. Davis and Daniel L. Davis. Verified by Travis J. LaDuc. Biodiversity Collections, University of Texas at Austin (TNHC 95867 [DRD 2198]). Two additional specimens were collected at this same locality on 8 August 2015 (TNHC 956865 [DRD 2172], 956865 [DRD 2173]). Numerous individuals have been observed at this location for the past two decades, representing a well-established population. New county record (Dixon 2013. *Amphibians and Reptiles of Texas: with Keys, Taxonomic Synopses, Bibliography, and Distribution Maps*. Texas A&M University Press, College Station, Texas. 447 pp.). Additional records of this species exist throughout Bastrop County (TNHC 66778, 66779, 85309). This introduced species has been documented in all surrounding counties (Caldwell, Fayette, Lee, Travis, Williamson) and throughout large regions of south, central, and northeastern Texas (Dixon 2013, *op. cit.*). Specimens collected under Texas Parks and Wildlife Scientific Collecting Permit (SPR-0511-126) issued to DRD.

DREW R. DAVIS, Department of Biology, University of South Dakota, 414 East Clark Street, Vermillion, South Dakota 57069, USA (e-mail: drew.davis@usd.edu); **DANIEL L. DAVIS**, Redfield Farm, 2158 FM 812, Dale, Texas 78616, USA.

MEROLES SQUAMULOSUS (Common Rough-scaled Lizard). SOUTH AFRICA: LIMPOPO PROVINCE: Lapalala Wilderness (23.906120°S, 28.306676°E; WGS 84). 15 March 2015. Simon Baeckens and Jan Scholliers. Verified by Werner Conradie. Virtual museum collection of the Southern African Reptile Conservation Assessment (SARCA No. 155809). One adult male was found on hard gravel terrain sprinting between vegetation patches. This large specimen (SVL = 69.14 mm) counted 15 femoral pores on each thigh. Here, we present the first record of this lacertid species for the Lapalala Wilderness nature reserve, and subsequently the first record for the SARCA quarter-degree grid 2328CD of approximately 676 km² (Bates et al. 2015. *The Reptile Atlas and Checklist of South Africa, Lesotho and Swaziland*. SANBI, Pretoria. 164 pp.). Fieldwork was conducted under a permit from the Limpopo Department of Economic Development, Environment and Tourism (No. 0092-MKT001-00004).

SIMON BAECKENS (e-mail: simon.baeckens@uantwerp.be), **JAN SCHOLLIERS** (e-mail: jan.scholliers@uantwerp.be), and **RAOUL VAN DAMME**, Department of Biology, Laboratory of Functional Morphology, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium (e-mail: raoul.vandamme@uantwerp.be).

OPHISAURUS ATTENUATUS (Slender Glass Lizard). USA: FLORIDA: MADISON Co.: Twin Rivers State Forest (30.26944°N, 83.26164°W; WGS 84). 27 April 2015. Jonathan D. Mays. Verified by Kenneth L. Krysko. Florida Museum of Natural History (UF 175552). Individual trapped by a drift fence installed in sandhill habitat. Specimen collected after it was depredated by fire ants. First voucher for the county (Krysko et al. 2011. *Atlas of Amphibians and Reptiles in Florida*. Final report, Florida Fish and Wildlife Conservation Commission, Tallahassee, Florida. 524 pp.).

E. PIERSON HILL (e-mail: pierson.hill@myfwc.com), **JONATHAN D. MAYS**, and **KEVIN M. ENGE**, Florida Fish and Wildlife Conservation Commission, 1105 S.W. Williston Road, Gainesville, Florida 32601, USA.

PLESTIODON EGREGIUS (Mole Skink). USA: FLORIDA: GULF Co.: 199 Beach Street, Port St. Joe (29.865934°N, 85.343002°W; WGS 84). 28 March 2010 E. Pierson Hill. Verified by Kenneth L. Krysko. Florida Museum of Natural History (UF177226, photo